

(X) R1 ( ) R2 ( ) R3 ( ) PG0 ( ) PG1 ( ) Estagiário ( ) Tecnólogo ( )  
PIBIC Last Name - Lobo First Name - Rodrigo Antonio Middle - Barbosa

Service (sector) Trauma N° CEP

### **Direct Cyclopexy for Traumatic Ciliochoroidal Detachment**

Lobo, Rodrigo MD; Magosso, Larissa MD; Mitne, Somaia MD; Paranhos, Augusto MD; Moraes, Nilva S.B. MD.

Purpose: To describe the direct cyclopexy for a traumatic ciliochoroidal detachment with persisting ocular hypotony.

Methods: A 41 year-old man complaining of progressive blurred vision in his left eye (OS) after ocular blunt trauma was referred to the emergency room of Sao Paulo Hospital (UNIFESP). His best-corrected visual acuity was hand motions and the ophthalmologic examination revealed severe ocular hypotony with a small cyclodialysis cleft. Indirect ophthalmoscopy showed severe hypotonic maculopathy. A peripheral choroidal detachment and a 360° ciliochoroidal detachment were found at the ultrasound examination and ultrasound biomicroscopy (UBM). Direct surgical cyclopexy was performed in 4 quadrants without intraoperative complications. Serial UBM exams and retinographs were performed during postoperative period.

Results: After 1 month follow-up intra-ocular pressure was controlled with no medication and at last visit (5 months after surgery) the visual acuity was 20/40 with no signs of maculopathy or any other ophthalmoscopy findings.

Conclusion: The ciliochoroidal detachment is a serious complication of severe ocular trauma and must be repaired before retinal folds lead to irreversible visual loss. Direct surgical cyclopexy is a successful treatment for a large hypotonous ciliary body detachment.